# SOLO® DENTAL NEEDLES SOLO NEEDLES: Sterile stainless steel siliconized dental needles with bevel marking

The needles are intended for maxillofacial injections. The needles are indicated for routine local dental anaesthesia injections.

## CONTRAINDICATION

Hypersensitivity to any component.

- INSTRUCTIONS FOR USE

  Break the sterlity guarantee seal just before use by turning the two parts of the protective sleeve in opposite directions. Screw the pre-threaded needle onto the pre-threaded hub of the syringe.
- Remove the plastic cap just before the injection is administered. It can be used again to remove the needle from the syringe to avoid risk of needle stick injury.
- Discard the used needle into a puncture-resistant container in occordance with all applicable with medical waste regulations.

(CE):12/1995

Rev. November 2019

- Never recap needles by hand. Use only accepted dental recapping procedures. Recapping needles by hand or recapping bent or broken needles can cause accidental needle stick injury.

- Avoid extreme pressure and excessive needle movement during injection as this may cause needles to break, resulting in serious injury to you and/or your patient.

Pug	Short	Cara Short	Solo	
0.4 × 35	0.3 × 23	0.3 x 12	@ (mm) x i. (mm)	
27 .	30	30	Ø (Ga)	
Nerve Block	Periopical	Intraligmentary	Type of Injection	

- Do not bend, break, or otherwise stress needles as serious injuries to you and/or your patient can occur.
- Do not insert needle to hub during injections, as needles can break and become lodged in patient's tissue, potentially causing serious permanent injury.
- Do not use short needles (<30 mm) when the expected depth in</li> soft tissue approaches the length of the needle.
- Be especially vigilant when positioning a needle in a patient who appears to be apprehensive.

# PRECAUTIONS FOR USE:

- Store in original packaging until ready to use.
- Read the Instructions for use supplied with your injection device.
- If seal is broken, the needle IS NOT STERILE and MUST NOT BE USED.
- For single use only: this device is for use in a single patient.
   Any reuse exposes other patients and the user to critical risks, including infection and trauma since the needle is no longer sterile and is more likely to break.
- If multiple injections are required, replace the needle for each cartridge in order to minimise the risk of needle point damage.
   1 patient / 1 cartridge / 1 needle
- The choice of the needle depends on the use by the practitioner.

### STORAGE

Do not store above 30°C and protect from maisture.

## PRESENTATION

100 two-point disposable non-resterillisable dental needles.