





How to use a Coloplast bedside drainage bag with Coloplast bag hanger or stand

Preparation



Simpla S

Bag hanger / stand

Conveen Basic Simpla



DC







1. A bedade drainage bag can be attached directly to a unsheath or indivelling catheter

Or it can be connected to a leg bag for extradrainage capacity during the night. For overnight linkage the leg bag must be fitted with a flexible outlet tube that will accommodate the inlet connector of the bodside drainage hag

Intermittent cathetensation

A bedside drainage bag may also be connected to an intermittent catheter for urinary collection during intermittent catheterisation





Instructions for use CM00000000407 Version 2

The Coloplast logo is a registered trademark of Coloplast A/S. All rights reserved ©






2. If the bag has a sample port, a sample of urine can be taken when necessary

Use a syringe with needle, or a needle free syringe as required.







When cutting a corrugated tube, leave as much of the straight section as possible in place. This will ensure a secure connection between tube and connector when mounting the connector, or between bag and tube when mounting the tube.

Indication

Urine collection bag

Information

Coloplast accepts no liability for any injury or loss that may arise if this product is used in a manner contrary to Coloplast's current recommendations

According to national guidelines it is not recommended to attach a non-sterile bedside drainage bag directly to an indwelling catheter.

Do not use if package is damaged.

Re-use of the single use product may create a potential harm to the user. Reprocessing, washing, disinfection, and/or sterilisation may compromise product characteristics, causing additional risk of physical harm to or infection of the user.





4. Attach the connector to the tube by pushing the thin end of the connector into the tube as far as possible.

Or attach the tube to the bag by pushing the cut end of the tube firmly into the connection inlet of the bag.





5. Use a Coloplast bag hanger or stand to suspend the bedside drainage bag.

Fit the bag securely on the hanger or stand by inserting the teeth of the hanger or stand in the slits at the top of the bag. Hook the bag hanger onto the bed rail or place the stand on the floor at the bedside.

To ensure optimal drainage, it is important that the bedside drainage bag is positioned lower than yourself when you are lying in bed.

Intermittent catheterisation Suspend or place the bag as you find most convenient, making sure the bag is positioned lower than your pelvic area whether you prefer to catheterise sitting, standing or lying down.











6. Make sure the outlet tap is closed.

Application





Attach the connector to the unsheath or indwelling catheter.

7. If the connector has a protective cap,





The urinary collecting system is now functional.



Intermittent catheterisation Attach the connector to the intermittent

When you begin to catheterise the urine will drain into the bedside drainage bag.





8. We recommend emptying the bag when it is a little over half full to ensure optimal drainage.

Intermittent cathetensation Empty the bedside drainage bag after each intermittent catheterisation.









Disconnect the bedside drainage bag from the urisheath or indwelling catheter.

Or, if the bedside drainage bag is linked to the leg bag, close the leg bag outlet tap. Then disconnect the bedside drainage bag. Avoid touching the open end of the leg bag outlet tube.

Intermittent catheterisation After completing the catheterisation, disconnect the bedside drainage bag from the intermittent catheter.

Detach the bag from the hanger or stand.



10. Change the bag according to local guidelines and recommendations.

The bag hanger or stand may be reused.

Empty the bag before discarding it.

The bag is intended for single use only and should be disposed of with normal household waste. Do not flush the product down the toilet.

Symbol



Do not use if package is damaged